

GALAXY FORMATION ON A MOVING MESH

Dušan Kereš

Hubble Fellow

Theoretical Astrophysics Center, UC Berkeley

Collaborators: Mark Vogelsberger, Lars Hernquist, Debora Šijački, Volker Springel

How to reliably simulate galaxies in cosmological environment?

- Formation of galaxies requires proper modeling of:
 - Structure formation
 - Gas infall into galaxies
 - Outflows out of galaxies
 - Interactions of infall/outflows/galaxies with the IGM/CGM
- Complex, nonlinear processes, large dynamical range!
- Successful codes applied to these problems need to:
 - Be adaptive to cover huge dynamical range.
 - Quickly and accurately calculate gravitational interactions.
 - Properly model hydrodynamics: discontinuities, steep gradients, shocks, instabilities, shear flows etc.
- Both SPH and AMR have weaknesses in some of these areas.

Some Strengths & Weaknesses

SPH


- accurate gravity solvers
- Galilean invariant
- spatially/temporally adaptive
- continuous refinement
- flexible geometries
- shocks broadened
- discontinuities not well-resolved
- relatively diffusive (artificial viscosity)
- instabilities suppressed
- limited mass resolution
- mixing suppressed

Eulerian-AMR


- accurate shock solvers (Godunov)
- resolution of discontinuities
- relatively less diffusive
- spatially/temporally adaptive
- large dynamic ranges
- mixing
- less accurate gravity solvers
- not Galilean invariant
- discontinuous refinement
- refinement criteria
- less flexible geometries

Mostly complementary attributes

Hybrid approach: moving mesh AREPO (Springel 2010)



- Voronoi tessellation of the computational domain
- Locations, motion of mesh-generating points arbitrary
- AREPO can mimic pure Lagrangian, static mesh & AMR codes
- If mesh-generating points move with fluid velocity: Galilean-Invariant
- Example: Kelvin-Helmholtz instability on 50 x 50 mesh




From V. Springel

Cosmological simulations with moving mesh and SPH


- Moving mesh code AREPO and SPH code GADGET-3.
- Start with the same ICs: $20/h$ Mpc box realized with 2×128^3 , 2×256^3 , 2×512^3 particles/cells:
 - Λ CDM cosmology, UV background, primordial cooling curve.
 - Star formation prescription in pressurized medium of Springel & Hernquist (2003), no outflows from galaxies.
 - Standard implementations, no parameter change to make results agree.
- AREPO is built on GADGET frame: uses the same gravity solver.
- AREPO in Lagrangian mode: relatively constant mass in a fluid element, set to be similar to SPH particle mass.
 - We add refinement and de-refinement, that are only occasionally applied to keep the mass in cell even closer to SPH mass over time.

Results on different scales


- *AREPO runtimes are only ~1.2-1.5 times slower than GADGET*



Extended disks, without ejective feedback...




Stellar disks are also more extended




Radii of cold galactic gas

Gaseous disks of galaxies are systematically more extended in AREPO.



SFR-history

After $z \sim 3$, AREPO runs have higher SFR density.




Higher SFRs in massive halos -> efficient hot gas cooling in AREPO


-Late time difference in global star formation is caused by massive halos

GADGET

AREPO



Massive halos in AREPO have higher central densities and lower central entropies



$z=0$

What causes the differences?

- Sizes and efficient gas cooling in massive galaxies are likely connected.
- AREPO:
 - Extended, gas rich disks are easily stripped in hot halos.
 - Stripped material will not lose angular momentum via dynamical friction like the rest of the infalling substructure (e.g. Maller&Dekel 2002)
 - It is efficiently mixed with hot gas, enabling more efficient cooling of hot halo.
- GADGET:
 - Harder to strip compact disks with lower gas fractions.
 - Clumps of gas form via cooling instability and from the stripped material.
 - These clumps survive too long owing to inability of SPH to properly capture hydro-instabilities (Agertz et al. 2007).
 - Clumps can heat the surrounding gas and lose angular momentum during the infall.
 - This process can lower cooling efficiency of the hot gas and cause transfer of angular momentum to the hot gas.
- Large differences in energy dissipation of sub-sonic turbulence in the halo infall region and differences in shock capturing create stronger heating of gas at intermediate radii in GADGET.

CONCLUSIONS

- AREPO is a very efficient code, suitable for cosmological simulations of galaxy formation.
- Gas in centers of hot halos in AREPO cools more efficiently than in the SPH code GADGET.
- Specific angular momentum of galactic gas is higher and more aligned in AREPO -> more extended, regular, gaseous disks.
- AREPO will likely improve our understanding of important aspects of galaxy and IGM evolution (e.g. disk formation, halo absorbers, hot mode accretion and more).

Mark Vogelsberger





Harvard-Smithsonian Center for Astrophysics
Institute for Theory and Computation



Transition from cold to hot halos

GADGET



AREPO

