

# Observations of satellite galaxies in groups at $0 < z < 2.5$

Tomer Tal

UC Santa Cruz

UCSC Galaxy Workshop

8/12/2013

NSF Postdoctoral Fellow

# Galaxy evolution in groups


Most galaxies reside and evolve in groups

- At low redshift (redshift surveys):
  - Individual halos
  - Overall environment and halo properties: statistically (e.g., clustering, lensing)

Alternative: statistical background subtraction


(does not require spec-z so can go to higher redshift)

# Statistical background subtraction






# Statistical background subtraction



# Cumulative number density


- Rank order does not evolve rapidly even in the presence of mergers
- Follow a galaxy population while **allowing galaxies to evolve in mass**

SDSS + NMBS




# Projected number density functions

- Lack of evolution
- **Remarkable balance between mergers and accretion**



# Projected number density functions


- Lack of evolution
- **Remarkable balance between mergers and accretion**
- Agreement with Guo+11 SAM



# Evolution in semi analytic models

## Insight from G11:

- Most satellites at  $1 < r/\text{Mpc} < 3$  are on extreme orbits
- Galaxies inside the virial radius gradually merge with primary






1. What is the potential for growth through mergers (is there enough mass to support observed growth)?
2. Are satellite galaxies even affected by their group environment?


# Typical massive galaxy environment

- Integrated number (mass) of satellites
- “Generalized” pair count measurement



# Typical massive galaxy environment

- Integrated number (mass) of satellites
- “Generalized” pair count measurement
- Equal total stellar mass in satellites as in primary




# Testing environmental effects

- Groups in a large redshift range from three surveys:
  - $z \sim 0$ : SDSS NYU-VAGC (Blanton+ 2005)
  - $z < 1.2$ : UltraVISTA (Muzzin+ 2013)
  - $z < 2.5$ : 3D-HST+CANDELS (Brammer+ 2012)
- Cumulative number density matching  $n=4 \times 10^{-4}$   
(corresponds to  $M(z=0) \sim 8 \times 10^{10} M_{\odot}$ )




# Rest frame UVJ

- Location on a UVJ diagram correlates well with star formation activity
- Simple cuts separate quiescent from star forming galaxies






# Primary/satellite selection






# Primary/satellite selection





# Evolution of the quiescent fraction

- Constant cumulative number density  $n = 4 \times 10^{-4}$
- All galaxy populations get quenched with time





# Summary

- Radial distribution of satellite galaxies in massive groups does not seem to evolve
- Remarkable balance between in-halo mergers and accretion of galaxies into the halo
- Star formation in group galaxies is affected by environmental processes